
REVIEW ARTICLE

The salience of existential concerns across the cancer control continuum

VIRGINIA LEE N., PH.D.,¹ AND CARMEN G. LOISELLE, N., PH.D.,^{2,3}

¹McGill University Health Centre, Montreal, Quebec, Canada

²School of Nursing and Department of Oncology, Faculty of Medicine, McGill University, Quebec, Canada

³Department of Psychosocial Oncology, Jewish General Hospital, Montreal, Quebec, Canada

(RECEIVED March 15, 2011; ACCEPTED April 9, 2011)

ABSTRACT

Despite significant improvement in cancer survival, the fear of death still remains rooted in individuals' beliefs about cancer. Existential fears pertaining to cancer cut across the cancer control continuum and taint decisions related to prevention, screening, surveillance, and follow-up recommendations, as well as the overall management of cancer-related issues. However, individuals are innately predisposed to cope with their cancer-related fears through mechanisms such as reliance on the process of meaning making. To better appreciate the potential impact of existential concerns across the cancer control continuum, the Temporal Existential Awareness and Meaning Making (TEAMM) model is proposed. This tripartite model depicts three types of perceived threats to life related to cancer including a "social awareness" (i.e., cancer signals death), "personalized awareness" (i.e., *I could die from cancer*), and the "lived experience" (i.e., *It feels like I am dying from cancer*). This construal aims to enhance our understanding of the personal and contextual resources that can be mobilized to manage existential concerns and optimize cancer control efforts. As such, existential discussions should be considered in any cancer-related supportive approach whether preventive, curative, or palliative, and not be deferred only until the advanced stages of cancer or at end of life. Further delineation and validation of the model is needed to explicitly recognize and depict how different levels of existential awareness might unfold as individuals grapple with a potential, actual, or recurrent cancer.

KEYWORDS: Cancer, Existential concerns, Meaning, Psychosocial adjustment, Cancer control continuum

INTRODUCTION

Despite significant gains in the relative 5-year survival rates for most cancers (Coleman et al., 2008; Canadian Cancer Society National Cancer Institute of Canada, 2010; Jemal et al., 2010), public perceptions about the survivability of cancer are less optimistic, often still being equated with death (Donovan et al., 2002, 2006; Holland et al., 2010). In fact, existential concerns generally lie latent in the general population (Becker, 1973; Neimeyer et al., 2004), and awareness about the fragility of life and the inevit-

ability of death only heighten when individuals are exposed to cancer-related cues (Benyamini et al., 2003; Klein & Stefanek, 2007; Lipworth et al., 2010).

Existential concerns are expected to influence behavior at any point of the cancer trajectory from relative adoption of illness-preventive actions, follow-up with cancer screening recommendations, and adherence to treatment modalities, to survivorship care. The purpose of this article is to capture the significance and ramifications of cancer-related existential issues across the cancer control continuum by proposing the Temporal Existential Awareness and Meaning Making (TEAMM) model. This tripartite model draws on key literature to delineate three distinct clinical presentations of existential awareness related to cancer. A discussion highlights the clinical

Address correspondence and reprint requests to: Virginia Lee, Nurse Scientist, McGill University Health Centre, 1650 Cedar Avenue Room S2-214, Montreal, QC H3G 1A4, Canada. E-mail: virginia.lee@muhc.mcgill.ca

and research implications of the TEAMM model for person-centered cancer care.

Distinguishing Between Existential Awareness and Existential Distress

A cancer diagnosis is often viewed as a pivotal life event, when the inevitability of one's own personal death becomes undeniably clear (Coyle, 2004). "Existential awareness" is the awareness of one's mortality and refers to what has been commonly called "mortality salience" in the literature (Burke et al., 2010). Existential awareness triggered by a cancer experience involves the contemplation of personal identity, autonomy, dignity, life meaning and purpose, and connections with others within the context of life and death (Frankl, 1959; Bolmsjo, 2000; Griffiths et al., 2002).

On the other hand, "existential distress" refers to what has been commonly called "death anxiety" (Pyszczynski et al., 1997) or "existential suffering" (Schuman-Olivier et al., 2008). The distress evoked by a cancer diagnosis often results from a subjective appraisal of a threat to one's life (Folkman et al., 1986) and not necessarily by more objective indices such as disease progression or prognosis (Cella & Tross, 1987; Laubmeier, 2004). Existential distress has been shown to be a separate construct from general anxiety, depression, somatic distress, and global psychological distress (Cella & Tross, 1987; Lichtenthal et al., 2009) and has been shown to correlate with time since diagnosis and closeness to death (Weisman & Worden, 1976–77; Cella & Tross, 1987; Lichtenthal et al., 2009).

Studies indicate that individuals with advanced cancer who have an accurate prognostic awareness are not necessarily distressed (Blinderman & Cherny, 2005; Barnett, 2006; Lichtenthal et al., 2009). Therefore, existential *awareness* does not necessarily lead to existential *distress* (Blinderman & Cherny, 2005). The appraisal of life threat from cancer can be construed on a continuum that ranges from a normative acceptance that is not necessarily upsetting to incrementally distressing levels characterized by ruminative thoughts and feelings of demoralization, meaninglessness, and isolation (Lee, 2008).

***Memento Mori*¹ and the Need for Meaning in the Context of Cancer**

Cancer is perhaps one of the most timeless representations of *memento mori* in health and illness (Dono-

van et al., 2002). Research into the existential unrest evoked by cancer was initially treated in the seminal work of Weisman and Worden (1976–77), which described a predominance of life and death concerns in the first 100 days following the diagnosis. Not surprisingly, existential concerns permeate the acute and survivorship stages of cancer. There is a preponderance of articles about the existential concerns at end of life and palliative care phases of the cancer trajectory. However, the literature is also suggesting the important influence of existential issues on cancer screening and preventive behaviors such as breast self-examination and mammography (Goldenberg et al., 2008, 2009), smoking cessation (Hansen et al., 2010), adherence to protective skin cancer behaviors (Cameron, 2008; Garside et al., 2010); uptake of fecal occult blood testing (Chapple et al., 2008), and genetic testing for cancer (Espalen, 2003).

Terror Management Theory (TMT) proposes that individuals have a basic instinctual drive for survival and self-preservation (Greenberg et al., 2000). When confronted with the idea of one's potentially imminent and inevitable death, individuals commonly rely on the use of *proximal defense mechanisms* (e.g., denial, distraction, avoidance, and cognitive distortion) or the use of *distal defense mechanisms* (e.g., efforts to maintain a sense of meaning, purpose, and permanence) to cope with the thought of one's potential nonexistence.

A burgeoning body of evidence exists to support the notion that many individuals search to reconstruct a sense of meaning in the face of life's tragedies (Frankl, 1959; Taylor, 1983; Thomson & Janigian, 1988; Park & Folkman, 1997; Breitbart, 2002; Lee et al., 2004, 2008; Park, 2010). Patients with life-altering cancer diagnoses and treatments often cannot look to a cure to relieve their anxieties and to restore equanimity (i.e., proximal defense mechanisms can be effective only up to a certain point). Instead, individuals with cancer often benefit from a new integration of the meaning of the illness into their lives. Efforts to facilitate this exploration of meaning or "meaning making" has recently been welcomed with a high degree of consensus by theorists, researchers, and clinicians who advocate for the importance of "meaning" in the context of cancer and other highly stressful life experiences including a potential, actual, or recurrent cancer diagnosis that challenges one's existence (Folkman & Greer, 2000; Park, 2010 for an extensive review). A broader appreciation of the existential issues across the cancer control continuum can inform the targeted development of meaning-oriented approaches to support health-protective lifestyle behaviors and early cancer detection, as well as actual experience with cancer.

¹*Memento mori* is a Latin phrase that names a genre of artistic creations whose main purpose is to remind people of their own mortality.

METHODS

A literature search pertaining to the existential discourse of the adult patient and cancer was conducted using the databases ISI Web of Knowledge, Medline, and PsychInfo for articles published in the English language between 1975 and 2010. The following search terms were used: “existential” in combination with “cancer,” “meaning making,” “death anxiety,” “terror management,” “risk,” “prevention,” “screening,” “genetics,” “end of life,” and “palliative.” Articles with a focus on children, adolescents, caregivers, family, and health professionals were excluded.

The cancer control continuum (National Cancer Institute, 2007; Miller et al., 2009) was selected to organize the predominant existential themes emerging from the literature search. The cancer control continuum was chosen for its broad definition of the cancer experience that begins earlier than at diagnosis, and for explicitly including the existential domain within the psychosocial issues that cut across different types of cancer diagnoses. The articles retrieved were closely read to identify (1) the foci and the gaps along the cancer control continuum in terms of existential discourse in cancer, and (2) the dominant themes emerging from the existing discourse.

RESULTS

Temporal Existential Awareness and Meaning Making (TEAMM) Model

The TEAMM model provides a guide to understanding how cancer-induced existential awareness can affect individuals across all phases of the cancer control continuum (National Cancer Institute, 2007; Miller et al., 2009). Three dominant themes were extrapolated from the literature to represent three types of cancer-related existential awareness: *a social awareness*, *a personalized awareness*, and *a lived experience of cancer’s threat to life* (Fig. 1). According to the literature, these three types of existential awareness are found to: (1) extend across *any* phase of the cancer control continuum (including individuals who are cancer free), (2) co-occur in varying intensities, and (3) be heightened or mitigated by one’s repertoire of coping strategies. Table 1 presents a clinical classification system to summarize the unique defining attributes for each type of cancer-related existential awareness. This system proposes that each type of existential awareness is differentially triggered by distinct cues, affects one’s sense of personal identity, and is managed differently through proximal or distal defense mechanisms. Figure 2 offers a pictorial representation of how individuals might confront, manage, and alternate




Fig. 1. Three types of existential awareness from cancer’s threat to life.

among the different types of existential awareness across each phase of the cancer control continuum. The TEAMM model is an extension of the study findings of Little and Sayer (2004), which describes how individuals are rendered “mortality salient” upon initial diagnosis of cancer, “*death salient*” when in remission, and “*dying salient*” when close to death.

A Social Awareness of Cancer’s Threat to Life (“Cancer Signals Death”)

The “cancer equals death” equation is a universal belief that is held even by individuals free of cancer (Borland et al., 1994; Donovan et al., 2002; Benyamini et al., 2003; Peters et al., 2006). Social awareness of death from cancer is often seen as a transient, unspoken acknowledgement of potential death from cancer described by experts in the field as the “elephant in the room” (Knapp-Oliver & Moyer, 2009; Holland et al., 2010; Lipworth et al., 2010). Educational and public health campaigns that aim to eradicate cancer or enhance the quality of life of people affected by cancer are examples of the explicit acknowledgement of cancer’s threat to life. Social expressions of cancer fears are also implicit in the stigma and shame that remains with certain types of cancers on the basis of anatomical location (Gray et al., 2000; Madlensky et al., 2004; Rosman, 2004; Chapple et al., 2008; Else-Quest et al., 2009) or perceived personal responsibility (Chapple et al., 2004; Bell et al., 2010; Gulyn & Youssef, 2010).

Research has shown that the strong negative affect elicited by cancer cues are transient because

Table 1. *The Temporal Existential Awareness and Meaning Making (TEAMM) model*

	Type of existential awareness		
	Social awareness	Personalized awareness	Lived experience
Cancer cues	Educational or public health campaigns; history of cancer (late diagnostics, lack of screening, stigma); personal experience with family and friends with cancer	Genetic testing; screening procedures leading to actual or potential personal life change; disruption in personal meaning; temporary or permanent inability to complete personal goals	Insurmountable distress from a single or cluster of physical or psychological symptoms
Predominant characteristics	Cognitive, affective	Cognitive, affective	Corporeal, cognitive, and affective
Integrity of self	Abstract, transient	Perceived threat	Under actual threat
Perception of personal vulnerability	Untouched	Present but threat of death is not imminent	Acute perception that death is imminent
In the absence of a personal history with cancer	Absent or transient	Heightens with perception of personal vulnerability and risk	May occur in the presence of a severe symptom experience
	Heightens with perception of increased personal vulnerability and risk	Diminishes in response to defensive coping strategies	Diminishes in response to defensive coping strategies and/or resolution of symptom
In the presence of a personal history with cancer	Diminishes in response to defensive coping strategies	Heightens when cancer enters into life context and/or symptoms become distressing	Heightens when physical or psychological symptoms are severe, uncontrolled, and/or unrelenting
	Heightens when symptom distress recedes or resolves	Diminishes when symptoms recede or resolve	Diminishes when symptoms recede or resolve
	Decreases relative to personalized awareness of death with progression of disease		
Coping	Collective and social	Individual and social	In isolation
Proximal defenses	Effective	Partially effective	Ineffective or maladaptive
Distal defenses	Unchallenged	Possibly challenged	Possibly challenged
	Intact	Search for meaning may be initiated	Search for meaning may be initiated

such cues activate protective coping mechanisms, namely, optimistic bias, denial, or avoidance, to suppress a rising awareness of the threats associated with cancer (Arndt et al., 1998, 2007; Benyamini et al., 2003; Lipworth et al., 2010). The use of proximal defence mechanisms that minimize risk perception and vulnerability, for example, have been reported even amongst individuals who have a family history of cancer (Benyamini et al., 2003; Madlensky et al., 2004). Other defensive behaviors include social

distancing from individuals with cancer (Mosher & Danoff-Burg, 2007) and avoidance or delay in seeking treatment (Tod & Joanne, 2010). When these coping mechanisms are activated, cancer is not explicitly appraised as personally life threatening. One's worldview remains relatively intact, and there is no perceived need to search for meaning (i.e., distal defense mechanisms are not affected) (Park & Folkman, 1997; Thomson & Janigian, 1988; Park, 2010).




Fig. 2. Predominant types of existential awareness across phases of the cancer control continuum.

A Personalized Awareness of Cancer's Threat to Life ("I Could Die from Cancer")

When cancer cues threaten the sense of order and purpose in one's own life context, cancer can no longer be perceived as an abstract illness that only happens to others. The individual enters into a personalized awareness of death from cancer. This awareness leads to a subjective appraisal of personal threat to one's integrity that does not necessarily correlate with objective signs of disease progression or prognosis (Blinderman & Cherny, 2005; Barnett, 2006; Lichtenthal et al., 2009). Individuals attempt to reconcile their own existence with the inevitable state of nonexistence upon death. The awareness of a personal life threat from cancer commonly occurs upon initial confirmation of a cancer diagnosis (Weisman & Worden, 1976–77; Derogatis, 1983; Landmark, 2001). However, a personalized awareness of death may also be evoked early in the absence of a personal history of cancer if significant distress is triggered by cancer cues related to screening procedures, genetic testing, or vicarious experiences involving others with cancer (Benyamini et al., 2003; Peters et al., 2006; Sanders et al., 2007). For others, the realization of one's own potential death from cancer may only develop later when the individual faces a progression, recurrence, or exacerbation of disease (Olsson et al., 2002; Richer & Ezer, 2002; Sarenmalm et al., 2007; McClement & Chochinov, 2008; Mehnert et al., 2009). Thus, a personalized awareness of death occurs when a cancer cue is appraised as personally threatening to one's survival.

In the absence of illness, the appraisal of life threat from cancer can be managed through the use of proximal defense mechanisms (Sanders et al., 2007; Lipworth et al., 2010). In the presence of actual disease, cancer progression, or debilitating side effects from treatment, it is hypothesized that the effectiveness of proximal or distal defenses may diminish as it becomes increasingly difficult to deny or minimize the impact of cancer on one's sense of integrity (Zimmermann, 2004, 2007; Zimmermann & Rodin, 2004). Individuals commonly cope by actively searching for a set of beliefs that can return a sense of order and permanence, as well as self-esteem and personal value, in the face of potential death from cancer (Taylor et al., 1984; O'Connor et al., 1990; Ruff Dirksen, 1995; Lee et al., 2004; Quinn, 2005; Kernan & Lepore, 2009).

The Lived Experience of Cancer's Threat to Life ("It Feels Like I Am Dying from Cancer")

The lived experience of death from cancer is posited as a corporeal, cognitive, and affective awareness of one's proximity to death. This transient sense of

"living while dying" is theorized to occur when the perceived severity of a single or a cluster of physical or psychological symptoms exceed one's resources and are considered insurmountable by the individual (Kissane & Clarke, 2001; Armstrong, 2003). The perception of proximity to death is theorized to diminish when patients experience symptom relief from treatment or when they receive effective palliative care (Knuti et al., 2003; Lundstrom et al., 2009). Therefore, the lived experience of death can occur at any point along the cancer continuum depending upon the appraised severity and cause of the symptoms. For example, when patients experience fatigue, anorexia, and weight loss, they may appraise this symptom cluster as worrisome in the pre-treatment phase but normalize it as an expected side effect in the active treatment phase. If the symptom cluster continues to be unrelenting and uncontrolled, the patient may interpret it as a sign that the disease is progressing and that death is approaching in the post-treatment phase (Poole & Froggatt, 2002; Lindqvist et al., 2004; Potter, 2004; McClement, 2005; Wainwright et al., 2007).

Individuals who are living cancer's threat to life cope by isolation and social withdrawal, and a characteristic "turning inwards" (Little & Sayers, 2004). For some, the "lived experience" may be the final impetus to begin the process of meaning-making as a coping strategy to alleviate the distress generated by awareness of one's imminent death. Past research has demonstrated that the use of avoidant-type coping styles may be associated with greater anxiety and distress (Watson et al., 1984; Carver, et al., 1993). Current therapeutic approaches with a strong existential orientation such as meaning-centered interventions are yielding significant benefits on standardized measures of psychosocial well-being, particularly for individuals at the end of life (Kissane et al., 2003; Chochinov et al., 2005a; Lee et al., 2006; Breitbart et al., 2010; Henry et al., 2010). Therefore, therapeutic interventions that facilitate the process of meaning-making during the cancer experience are providing early evidence that the reformulation of distal defense mechanisms protects against death anxiety.

DISCUSSION

Balancing the Will to Live and the Fear of Death

The struggle to understand how one can continue to live knowing that death is imminent from cancer is a common paradox reported by many individuals with cancer (Halstead & Hull, 2001) and can be explained by the co-occurrence of the three types of awareness.

When different states of death awareness occur simultaneously, the focus on self-preservation and self-integrity is expected to predominate. Therefore, as depicted in the different scenarios in Figure 2, the “lived experience” of cancer’s threat to life is hypothesized to trump a “personalized awareness” of death from cancer. Similarly, in the absence of a corporeal experience of cancer, a “personalized” awareness of death from cancer is likely to compete with or overshadow a “social awareness” of cancer’s threat to life. Alternatively, the “social awareness” of cancer’s threat to life is hypothesized to regain strength and importance as symptoms resolve and individuals refocus their efforts on goals other than their own survival. These experiences do not necessarily follow a straightforward trajectory, as illustrated in Figure 2. The notion of relative intensity depicted by the three types of awareness is corroborated by a number of rich, descriptive studies in which people may feel as if they are swinging back and forth on a pendulum (Halstead & Hull, 2001; Giske & Artinian, 2008; Sand et al., 2009).

Impact on Cancer Control Efforts

The TEAMM model outlines how existential concerns (conscious or unconscious) can influence the cancer symptom experience, and shape behaviors related to the perception of cancer risk and adjustment to the cancer experience. The ability to manage one’s existential concerns related to the fear of cancer directly influences how patients make treatment decisions, including whether to adopt, adhere, delay, or forego uncomfortable or invasive procedures for cancer control (de Nooijer et al., 2003). Existential concerns influence and are influenced by the interpretation of symptoms (Armstrong, 2003; Lindqvist et al., 2004; Lundstrom et al., 2009).

Emotions, fears, beliefs, and other personally relevant factors often come into play during the illness experience (Bekker, 2010). Research shows that individuals respond selectively to cancer cues: some may be fatalistic in their interpretation of perceived personal risk, whereas others are not (Hurley et al., 2002). Individuals with a high fear of cancer may interpret ambiguous symptoms with a negative bias (Miles et al., 2009). They may be more motivated to engage in behaviors to protect one’s integrity and survival, whether by adhering to cancer screening practices or by actively searching for meaning (Andersen & Cacioppo, 1995). For example, researchers (Audrain et al., 1995; Lerman et al., 1995) have demonstrated that women were more motivated to undergo mammography when they held inaccurate and biased views of cancer risk that heightened their fear of death. For other individuals, new and unex-

pected symptoms are tempered by an optimistic bias to block the unpleasant thought of one’s potential for death by minimizing individual risk perceptions or reframing them as transient, self-correcting conditions (Andersen & Cacioppo, 1995). Aversion and low adherence rates to certain cancer screening procedures or behaviors that involve deliberate and explicit bodily manipulation and /or physical discomfort such as breast self examinations, mammography, fecal occult blood testing, or sigmoidoscopy have been linked to reminders of one’s “creatureliness” (Madlensky et al., 2004; Chapple et al., 2008; Goldenberg et al., 2008, 2009; Garside et al., 2010). Of particular interest is the work of researchers who document that explicit messages warning of the link between cancer and death may have the unintended opposite effect of increasing willingness to continue smoking particularly among individuals whose self-esteem is enhanced by smoking (Hansen et al., 2010). Clearly, individuals’ motivations to comply, delay, or reject cancer prevention practices is complex and related to existential concerns. The challenge for clinicians and public health professionals lies in marketing cancer control information in ways that invoke optimal levels of existential awareness so that individuals are motivated to undergo screening tests and behavior change (Klein & Stefanek, 2007).

Promotion of Existential Awareness

The evidence thus far indicates that once initiated, existential awareness persists over time. However, this awareness may vary in intensity, depending upon symptom severity, disease progression, and the experience of hope and meaning in life. For some, the salience of one’s own mortality coupled with symptom distress and an uncertain view of the future, can be distressing and immobilizing, and can even influence the desire for hastened death (Breitbart et al., 2000; McClain et al., 2003; Chochinov et al., 2005b). For others, awareness of one’s mortality may have adaptive value by dramatically changing one’s life priorities to initiate the phenomenon of post-traumatic growth (Cordova et al., 2001; Tedeschi & Calhoun, 2004; Bellizzi & Blank, 2006; Holland & Weiss, 2008; Jim & Jacobsen, 2008; Park, 2008; Park et al., 2008). Thus, the paradox of death awareness lies in its potential to be both psychologically paralyzing and instrumental in mobilizing a tenacious will to live. Several studies demonstrate that individuals with cancer have unmet existential needs, are not distressed by existentially oriented discussions, and would like to have more existential discussions with their treating team (Moadel et al., 1999; Blinderman & Cherny, 2005; Lichtenthal, 2009).

The challenge in cancer control efforts lies in developing theory-based approaches to capitalize on the potential functional value of existential awareness. A number of novel evidence-based interventions that buffer against existential distress are available from the literature in the fields of mental health and palliative care (Kissane et al., 2003; Chochinov et al., 2005a; Lee et al., 2006; Breitbart et al., 2010; Henry et al., 2010). Innovative interventions can evolve from these existing approaches to promote clinically appealing cancer control interventions. Clinicians can use the TEAMM model to understand when each type of existential awareness is most salient and facilitate the use of coping strategies most appropriate to that specific type of awareness (Little & Sayers, 2004; Klein & Stefanek, 2007). The literature suggests that most individuals have the psychological resilience to cope with the awareness of death that cancer evokes. This means that existential discussions need not be delayed until the advanced stages of cancer or at end of life, but should take place when the need arises.

Existential discussions in the context of cancer control and cancer care need not be intense, philosophical, or time consuming (Blinderman & Cherny, 2005; Kvåle, 2007). When these discussions are introduced in a non-threatening, secular manner, they can become an integral part of many comprehensive cancer control interventions. Existential exchanges communicate a sense of value and respect for the individual, and can build trusting relationships that can influence subsequent critical health decisions. Timely referrals for assessment and services can be provided when the patient's needs exceed the clinician's resources, capability, or personal comfort (Holland & Reznik, 2005; Surbone et al., 2010).

The proposed TEAMM model offers a starting point to broaden our understanding of how patients respond to the threat to life across the pre-disease and disease phases of the cancer control continuum. We raise four areas amenable to further research: (1) further validation of the three types of existential awareness proposed in the model through in-depth qualitative exploration with case series or longitudinal designs, (2) further exploration of the temporal relationship between existential awareness and symptom distress (i.e., how do existential issues influence and how are they influenced by the symptoms experience?), (3) exploration of the cross-cultural relevance of the TEAMM model, and (4) development and evaluation of novel meaning-oriented interventions aimed at encouraging uptake of cancer screening and preventive practices.

CONCLUSIONS

The TEAMM model relies on a broad spectrum of scientific knowledge to propose an evolving understanding of existential issues as they apply to person-centered care, and lays the foundation for further exploration, research, and discussion. Models of cancer control pertaining to existential distress have not previously included the pre-disease or early disease phases (Nolan & Mock, 2004; Knight & Emanuel, 2007; Schuman-Olivier et al., 2008). Existential discussions are pertinent across the entire cancer control spectrum and do not need to be deferred until the advanced stages of cancer or at end of life. The TEAMM model contributes to the notion of the cancer control continuum by furthering understanding about the role of existential awareness even in the earlier, disease-free, phases of the cancer control continuum.

Rowland and Baker (2005) eloquently state that "Being disease free does not mean being free of the disease." This idea captures the long-lasting and pervasive impact of cancer (potential or actual) and its diverse effects on affected individuals and families across the life span. This article extends this proposition to suggest that existential issues evoked by cancer, an illness that epitomizes death, can be as powerful and pervasive in the pre-disease and early disease phases as they are in the survivorship and end-of-life phases. It is our hope that if the existential fear of dying from cancer is directly and sensitively addressed with knowledge, compassion, and care, that this, in turn, will help improve patient-provider relationships, enhance psychosocial adjustment, and reduce the distress underlying cancer and related communication problems, to optimize cancer control and cancer care interventions.

ACKNOWLEDGMENTS

Virginia Lee was supported by a postdoctoral research fellowship from the Canadian Institutes of Health Research (CIHR)-Psychosocial Oncology Research Training Program. The authors gratefully acknowledge Janet Childerhose for her helpful comments on earlier versions of the manuscript and the Neuromedia Services at the Montreal Neurological Institute and Hospital for formatting the figures.

REFERENCES

- Andersen, B.L. & Cacioppo, J.T. (1995). Delay in seeking a cancer diagnosis: delay stages and psychophysiological comparison processes. *British Journal of Social Psychology*, *34*, 33–52.
- Armstrong, T.S. (2003). Symptoms experience: A concept analysis. *Oncology Nursing Forum*, *30*, 601–606.
- Arndt, J., Cook, A., Goldenberg, J.L., et al. (2007). Cancer and the threat of death: The cognitive dynamics of

- death-thought suppression and its impact on behavioral health intentions. *Journal of Personality and Social Psychology*, 92, 12–29.
- Arndt, J., Greenberg, J., Simon, L., et al. (1998). Terror management and self-awareness: Evidence that mortality salience provokes avoidance of the self-focused state. *Personality and Social Psychology Bulletin*, 24, 1216–1227.
- Audrain, J., Lerman, C., Rimer, B., et al. (1995). Awareness of heightened breast cancer risk among first degree relatives of recently diagnosed breast cancer patients. *Cancer Epidemiology, Biomarkers, & Prevention*, 4, 561–565.
- Barnett, M.M. (2006). Does it hurt to know the worst? Psychological morbidity, information preferences and understanding of prognosis in patients with advanced cancer. *Psycho-Oncology*, 15, 44–55.
- Becker, E. (1973). *The Denial of Death*. New York: Collier–Mac.
- Bekker, H.L. (2010). The loss of reason in patient decision aid research: Do checklists damage the quality of informed choice interventions? *Patient Education and Counseling*, 78, 357–364.
- Bell, K., Salmon, A., Bowers, M., et al. (2010). Smoking, stigma and tobacco ‘denormalization’: Further reflections on the use of stigma as a public health tool. A commentary on *Social Science & Medicine’s* Stigma, Prejudice, Discrimination and Health Special Issue (67:3). *Social Science & Medicine*, 70, 795–799.
- Bellizzi, K.M. & Blank, T.O. (2006). Predicting posttraumatic growth in breast cancer survivors. *Health Psychology*, 25, 47–56.
- Benyamini, Y., McClain, C.S., Leventhal, E.A., et al. (2003). Living with the worry of cancer: Health perceptions and behaviors of elderly people with self, vicarious, or no history of cancer. *Psychooncology*, 12, 161–172.
- Blinderman, C.D. & Cherny, N.I. (2005). Existential issues do not necessarily result in existential suffering: lessons from cancer patients in Israel. *Palliative Medicine*, 19, 371–380.
- Bolmsjo, I. (2000). Existential issues in palliative care — Interviews with cancer patients. *Journal of Palliative Care*, 16, 20–24.
- Borland, R., Donaghue, N. & Hill, D. (1994). Illnesses that Australians most feared in 1986 and 1993. *Australian Journal of Public Health*, 18, 366–369.
- Breitbart, W. (2002). Spirituality and meaning in supportive care: Spirituality- and meaning-centered group psychotherapy interventions in advanced cancer. *Supportive Cancer Care*, 10, 272–280.
- Breitbart, W., Rosenfeld, B., Gibson, C., et al. (2010). Meaning-centered group psychotherapy for patients with advanced cancer: a pilot randomized controlled trial. *Psycho-Oncology*, 19, 21–28.
- Breitbart, W., Rosenfeld, B., Pessin, H., et al. (2000). Depression, hopelessness, and desire for hastened death in terminally ill cancer patients. *Journal of the American Medical Association*, 284, 2907–2911.
- Burke, B.L., Martens, A. & Faucher, E.H. (2010). Two decades of terror management theory: A meta-analysis of mortality salience research. *Personality and Social Psychology Review*, 14, 155–195.
- Cameron, L. (2008). Illness risk representations and motivations to engage in protective behaviour: The case of skin cancer risk. *Psychology & Health*, 23, 91–112.
- Canadian Cancer Society’s Steering Committee on Cancer Statistics. (2010). *Canadian Cancer Statistics 2011*. Toronto, ON: Canadian Cancer Society.
- Carver, C.S., Pozo, C., Harris, S.D., et al. (1993). How coping mediates the effects of optimism on distress—A study of women with early-stage breast cancer. *Journal of Personality and Social Psychology*, 65, 375–390.
- Cella, D.F. & Tross, S. (1987). Death anxiety in cancer survival- a preliminary cross-validation study. *Journal of Personality Assessment*, 51, 451–461.
- Chapple, A., Ziebland, S., Hewitson, P., et al. (2008). What affects the uptake of screening for bowel cancer using a faecal occult blood test (FOBT): A qualitative study. *Social Science & Medicine*, 66, 2425–2435.
- Chapple, A., Ziebland, S. & McPherson, A. (2004). Stigma, shame, and blame experienced by patients with lung cancer: qualitative study. *British Medical Journal*, 328, 1470.
- Chochinov, H.M., Hack, T., Hassard, T., et al. (2005a). Dignity therapy: A novel psychotherapeutic intervention for patients near the end of life. *Journal of Clinical Oncology*, 23, 5520–5525.
- Chochinov, H.M., Hack, T., Hassard, T., et al. (2005b). Understanding the will to live in patients nearing death. *Psychosomatics*, 46, 7–10.
- Coleman, M.P., Quaresma, M., Berrino, F., et al. (2008). Cancer survival in five continents: A worldwide population-based study (CONCORD). *Lancet Oncology*, 9, 730–756.
- Cordova, M.J., Cunningham, L.L.C., Carlson, C.R., et al. (2001). Posttraumatic growth following breast cancer: A controlled comparison study. *Health Psychology*, 20, 176–185.
- Coyle, N. (2004). The existential slap—A crisis of disclosure. *International Journal of Palliative Nursing*, 10, 520.
- de Nooijer, J., Lechner, L. & de Vries, H. (2003). Social psychological correlates of paying attention to cancer symptoms and seeking medical help. *Social Science & Medicine*, 56, 915–920.
- Derogatis, L.R., Morrow, G.R., Fetting, J., et al. (1983). The prevalence of psychiatric disorders among cancer patients. *Journal of the American Medical Association*, 249, 751–757.
- Donovan, J., Jalleh, G. & Jones, S. (2002). The word ‘cancer’: Reframing the context to reduce anxiety arousal. *Australian and New Zealand Journal of Public Health*, 27, 291–293.
- Donovan, R.J., Carter, O.B. & Byrne, M.J. (2006). People’s perceptions of cancer survivability: implications for oncologists. *Lancet Oncology*, 7, 668–675.
- Else-Quest, N.M., LoConte, N.K., Schiller, J.H., et al. (2009). Perceived stigma, self-blame, and adjustment among lung, breast and prostate cancer patients. *Psychology & Health*, 24, 949–964.
- Esplen, M.J. (2003). Adjusting to potentially life altering genetic information: Research findings on the role of existential group psychotherapy. *International Psychogeriatrics*, 15, 3–4.
- Folkman, S., Lazarus, R.S., Gruen, R.J., et al. (1986). Appraisal, coping, health status, and psychological symptoms. *Journal of Personality and Social Psychology*, 50, 571–579.
- Frankl, V. (1959). *Man’s Search for Meaning*, 4th ed. Boston: Beacon Press.
- Garside, R., Pearson, M. & Moxham, T. (2010). What influences the uptake of information to prevent skin cancer? A systematic review and synthesis of qualitative research. *Health Education Research*, 25, 162–182.

- Giske, T. & Artinian, B. (2008). Patterns of 'balancing between hope and despair' in the diagnostic phase: A grounded theory study of patients on a gastroenterology ward. *Journal of Advanced Nursing*, 62, 22–31.
- Goldenberg, J.L., Arndt, J., Hart, J., et al. (2008). Uncovering an existential barrier to breast self-exam behavior. *Journal of Experimental Social Psychology*, 44, 260–274.
- Goldenberg, J.L., Routledge, C. & Arndt, J. (2009). Mammograms and the management of existential discomfort: Threats associated with the physicality of the body and neuroticism. *Psychology & Health*, 24, 563–581.
- Gray, R.E., Fitch, M., Phillips, C., et al. (2000). To tell or not to tell: Patterns of disclosure among men with prostate cancer. *Psycho-Oncology*, 9, 273–282.
- Greenberg, J., Arndt, J., Simon, L., et al. (2000). Proximal and distal defenses in response to reminders of one's mortality: Evidence of a temporal sequence. *Personality and Social Psychology Bulletin*, 26, 91–99.
- Griffiths, C., Norton, L., Wagstaff, G., et al. (2002). Existential Concerns in late stage cancer. *European Journal of Oncology Nursing*, 6, 243–246.
- Gulyn, L.M. & Youssef, F. (2010). Attribution of blame for breast and lung cancers in women. *Journal of Psychosocial Oncology*, 28, 291–301.
- Hansen, J., Winzeler, S. & Topolinski, S. (2010). When death makes you smoke: A terror management perspective on the effectiveness of cigarette pack warnings. *Journal of Experimental Social Psychology*, 46, 226–228.
- Henry, M., Cohen, S.R., Lee, V., et al. (2010). The Meaning-Making intervention (MMi) appears to increase meaning in life in advanced ovarian cancer: A randomized controlled pilot study. <http://dx.doi.org/10.1002/pon.1764>.
- Holland, J. & Weiss, T. (2008). The new standard of quality cancer care: Integrating the psychosocial aspects in routine cancer from diagnosis through survivorship. *Cancer Journal*, 14, 425–428.
- Holland, J.C., Kelly, B.J. & Weinberger, M.I. (2010). Why psychosocial care is difficult to integrate into routine cancer care: Stigma is the elephant in the room. *Journal of the National Comprehensive Cancer Network*, 8, 362–366.
- Holland, J.G. & Reznik, I. (2005). Pathways for psychosocial care of cancer survivors. *Cancer*, 104, 2624–2637.
- Hurley, K., Du Hamel, K., Vickberg, S., et al. (2002). Does cancer-specific anxiety facilitate or deter screening? Different answers from breast and colon cancer screening. *Psychosomatic Medicine*, 64, 285.
- Jemal, A., Siegel, R., Xu, J., et al. (2010). Cancer Statistics, 2010. *CA: A Cancer Journal for Clinicians*, 60, 277–300.
- Jim, H.S. & Jacobsen, P.B. (2008). Posttraumatic stress and posttraumatic growth in cancer survivorship: A review. *The Cancer Journal*, 14, 414–419.
- Kernan, W.D. & Lepore, S.J. (2009). Searching for and making meaning after breast cancer: Prevalence, patterns, and negative affect. *Social Science & Medicine*, 68, 1176–1182.
- Kissane, D. & Clarke, D.M. (2001). Demoralization Syndrome—A relevant psychiatric diagnosis for palliative care. *Journal of Palliative Care*, 17, 12–21.
- Kissane, D.W., Bloch, S., Smith, G.C., et al. (2003). Cognitive-existential group psychotherapy for women with primary breast cancer: A randomised controlled trial. *Psycho-Oncology*, 12, 532–546.
- Klein, W.M. & Stefanek, M.E. (2007). Cancer risk elicitation and communication: Lessons from the psychology of risk perception. *CA: A Cancer Journal for Clinicians*, 57, 147–167.
- Knapp-Oliver, S. & Moyer, A. (2009). Visibility and the stigmatization of cancer: Context matters. *Journal of Applied Social Psychology*, 39, 2798–2808.
- Knight, S.J. & Emanuel, L. (2007). Processes of adjustment to end-of-life losses: A reintegration model. *Journal of Palliative Medicine*, 10, 1190–1198.
- Knuti, K.A., Wharton, R.H., Wharton, K.L., et al. (2003). Living as a cancer surprise: A doctor tells his story. *Oncologist*, 8, 108–122.
- Kvåle, K. (2007). Do cancer patients always want to talk about difficult emotions? A qualitative study of cancer inpatients communication needs. *European Journal of Oncology Nursing*, 11, 320–327.
- Landmark, B.T., Strandmark, M. & Wahl, A.K. (2001). Living with newly diagnosed breast cancer—The meaning of existential issues: A qualitative study of 10 women with newly diagnosed breast cancer, based on grounded theory. *Cancer Nursing*, 24, 220–226.
- Laubmeier, K.K. & Zakowski, S.G. (2004). The role of objective versus perceived life threat in the psychological adjustment to cancer. *Psychology and Health*, 19, 425–437.
- Lee, V. (2008). The existential plight of cancer: Meaning making as a concrete approach to the intangible search for meaning. *Supportive Care in Cancer*, 16, 779–785.
- Lee, V., Cohen, S.R., Edgar, L., et al. (2006). Meaning-making intervention during breast or colorectal cancer treatment improves self-esteem, optimism, and self-efficacy. *Social Science & Medicine*, 62, 3133–3145.
- Lee, V., Cohen, S.R., Edgar, L., et al. (2004). Clarifying 'meaning' in the context of cancer research: A systematic literature review. *Palliative and Supportive Care*, 2, 291–303.
- Lerman, C., Seay, J., Balshem, A., et al. (1995). Interest in genetic testing among first degree relatives of breast cancer patients. *American Journal of Medical Genetics*, 57, 385–392.
- Lichtenthal, W.G., Nilsson, M., Zhang, B., et al. (2009). Do rates of mental disorders and existential distress among advanced stage cancer patients increase as death approaches? *Psycho-Oncology*, 18, 50–61.
- Lindqvist, O., Widmark, A. & Rasmussen, B.H. (2004). Meanings of the phenomenon of fatigue as narrated by 4 patients with cancer in palliative care. *Cancer Nursing*, 27, 237–243.
- Lipworth, W.L., Davey, H.M., Carter, S.M., et al. (2010). Beliefs and beyond: what can we learn from qualitative studies of lay people's understandings of cancer risk? *Health Expectations*, 13, 113–124.
- Little, M. & Sayers, E.J. (2004). The skull beneath the skin: Cancer survival and awareness of death. *Psycho-Oncology*, 13, 190–198.
- Lundstrom, S., Furst, C.J., Friedrichsen, M., et al. (2009). The existential impact of starting corticosteroid treatment as symptom control in advanced metastatic cancer. *Palliative Medicine*, 23, 165–170.
- Madlensky, L., Esplen, M.J. & Goel, V. (2004). Reasons given by relatives of colorectal cancer patients for not undergoing screening. *Preventive Medicine*, 39, 643–648.

- McClain, C.S., Rosenfeld, B. & Breitbart, W. (2003). Effect of spiritual well-being on end-of-life despair in terminally-ill cancer patients. *The Lancet*, *361*, 1603–1607.
- McClement, S. (2005). Cancer anorexia–cachexia syndrome—Psychological effect on the patient and family. *Journal of Wound Ostomy and Continence Nursing*, *32*, 264–268.
- McClement, S.E. & Chochinov, H.M. (2008). Hope in advanced cancer patients. *European Journal of Cancer*, *44*, 1169–1174.
- Mehnert, A., Berg, P., Henrich, G., et al. (2009). Fear of cancer progression and cancer-related intrusive cognitions in breast cancer survivors. *Psycho-Oncology*, *18*, 1273–1280.
- Miles, A., Voorwinden, S., Mathews, A., et al. (2009). Cancer fear and the interpretation of ambiguous information related to cancer. *Cognition & Emotion*, *23*, 701–713.
- Miller, S.M., Bowen, D.J., Croyle, R.T., et al. (2009). *Handbook of Cancer Control and Behavioural Science. A Resource for Researchers, Practitioners, and Policymakers*. Washington, DC: American Psychological Association.
- Moadel, A., Morgan, C., Fatone, A., et al. (1999). Seeking meaning and hope: Self-reported spiritual and existential needs among an ethnically-diverse cancer patient population. *Psycho-Oncology*, *8*, 378–385.
- Mosher, C.E. & Danoff-Burg, S. (2007). Death anxiety and cancer-related stigma: A terror management analysis. *Death Studies*, *31*, 885–907.
- National Cancer Institute. (2007). *The Cancer Control Continuum 2007*. <http://cancercontrol.cancer.gov/OD/continuum.html>.
- Neimeyer, R.A., Wittkowski, J. & Moser, R.P. (2004). Psychological research on death attitudes: An overview and evaluation. *Death Studies*, *28*, 309–340.
- Nolan, M.T. & Mock, V. (2004). A conceptual framework for end-of-life care: A reconsideration of factors influencing the integrity of the human person. *Journal of Professional Nursing*, *20*, 351–360.
- O'Connor, A.P., Wicker, C.A. & Germino, B.B. (1990). Understanding the cancer patient's search for meaning. *Cancer Nursing*, *13*, 167–175.
- Olsson, U., Bergbom, I. & Bosaeus, I. (2002). Patient's experiences of the recovery period 3 months after gastrointestinal cancer surgery. *European Journal of Cancer Care*, *11*, 51–60.
- Park, C. (2008). Meaning and personal growth in the context of cancer. *International Journal of Psychology*, *43*(3–4), 752.
- Park, C. (2010). Making sense of the meaning literature: An integrative review of meaning making and its effects on adjustment to stressful life events. *Psychological Bulletin*, *136*, 257–301.
- Park, C., Blank, T. & Edmondson, D. (2008). Determinants and meaning of growth in cancer survivorship. *International Journal of Psychology*, *43*, 753.
- Park, C.L. & Folkman, S. (1997). Meaning in the context of stress and coping. *Review of General Psychology*, *1*, 115–144.
- Peters, E., McCaul, K.D., Stefanek, M., et al. (2006). A heuristics approach to understanding cancer risk perception: Contributions from judgment and decision-making research. *Annals of Behavioral Medicine*, *31*, 45–52.
- Poole, K. & Froggatt, K. (2002). Loss of weight and loss of appetite in advanced cancer: a problem for the patient, the carer, or the health professional? *Palliative Medicine*, *16*, 499–506.
- Potter, J. (2004). Fatigue experience in advanced cancer: A phenomenological approach. *International Journal of Palliative Nursing*, *10*, 15–23.
- Pyszczynski, T., Greenberg, J. & Solomon, S. (1997). Why do we need what we need? A terror management perspective on the roots of human social motivation. *Psychological Inquiry*, *8*, 1–20.
- Quinn, B. (2005). Cancer and the treatment: Does it make sense to patients? *Hematology*, *10*, 325–328.
- Richer, M.C. & Ezer, H. (2002). Living in it, living with it, and moving on: Dimensions of meaning during chemotherapy. *Oncology Nursing Forum*, *29*, 113–119.
- Rosman, S. (2004). Cancer and stigma: experience of patients with chemotherapy-induced alopecia. *Patient Education and Counseling*, *52*, 333–339.
- Rowland, J.H. & Baker, F. (2005). Introduction: Resilience of cancer survivors across the lifespan. *Cancer*, *104*, 2543–2548.
- Ruff Dirksen, S. (1995). Search for meaning in long-term cancer survivors. *Journal of Advanced Nursing*, *21*, 628–633.
- Sand, L., Olsson, M. & Strang, P. (2009). Coping strategies in the presence of one's own impending death from cancer. *Journal of Pain and Symptom Management*, *37*, 13–22.
- Sanders, T., Campbell, R., Donovan, J., et al. (2007). Narrative accounts of hereditary risk: Knowledge about family history, lay theories of disease, and “internal” and “external” causation. *Qualitative Health Research*, *17*, 510–520.
- Sarenmalm, E.K., Ohlen, J., Jonsson, T., et al. (2007). Coping with recurrent breast cancer: Predictors of distressing symptoms and health-related quality of life. *Journal of Pain and Symptom Management*, *34*, 24–39.
- Schuman-Olivier, Z., Brendel, D.H., Forstein, M., et al. (2008). The use of palliative sedation for existential distress: A psychiatric perspective. *Harvard Review of Psychiatry*, *16*, 339–351.
- Surbone, A., Baider, L., Weitzman, T.S., et al. (2010). Psychosocial care for patients and their families is integral to supportive care in cancer: MASCC position statement. *Supportive Care in Cancer*, *18*, 255–263.
- Taylor, S.E. (1983). Adjustment to threatening events: A theory of cognitive adaptation. *American Psychologist*, *38*, 1161–1173.
- Taylor, S.E., Lichtman, R.R. & Wood, J.V. (1984). Attributions, beliefs about control, and adjustment to breast cancer. *Journal of Personality and Social Psychology*, *46*, 489–502.
- Tedeschi, R.G. & Calhoun, L.G. (2004). Posttraumatic growth: Conceptual foundations and empirical evidence. *Psychological Inquiry*, *15*, 1–18.
- Thomson, S.C. & Janigan, A.S. (1988). Life Schemes: A framework for understanding the search for meaning. *Journal of Social and Clinical Psychology*, *7*, 260–280.
- Tod, A.M. & Joanne, R. (2010). Overcoming delay in the diagnosis of lung cancer: A qualitative study. *Nursing Standard*, *24*, 35–43.
- Wainwright, D., Donovan, J.L., Kavadas, V., et al. (2007). Remapping the body: Learning to eat again after

- surgery for esophageal cancer. *Qualitative Health Research*, 17, 759–771.
- Watson, M., Greer, S., Blake, S., et al. (1984). Reaction to a diagnosis of breast cancer-Relationship between denial, delay, and rates of psychological morbidity. *Cancer*, 53, 2008–2012.
- Weisman, A.D. & Worden, J.W. (1976). The existential plight in cancer: Significance of the first 100 days. *International Journal of Psychiatry in Medicine*, 7, 1–15.
- Zimmermann, C. (2004). Denial of impending death: A discourse analysis of the palliative care literature. *Social Science and Medicine*, 59, 1769–1780.
- Zimmermann, C. (2007). Death denial: Obstacle or instrument for palliative care? An analysis of clinical literature. *Sociology of Health and Illness*, 29, 297–314.
- Zimmermann, C. & Rodin, G. (2004). The denial of death thesis: Sociological critique and implications for palliative care. *Palliative Medicine*, 18, 121–128.